WETLAND DETERMINATION DATA FORM - Great Plains Region

L3R Project/Site: City	Marsha //County:	II	Sampling Date:	2015-06-06	
Enbridge Applicant/Owner:		Min State:	nesota	Sampling Point:	u-156n46w27-a1
ACM/KRG Investigator(s):	9	Section, Townsl	nip, Range:		
talf Landform (hillslope, terrace, etc.):			(concave, conv		0-2 Slope (%):
Subregion (LRR or MLRA):	Latitude:	48.303845613	3 Longit	-96.56568950 :ude:	
Minnesota State Plane North, NAD 83					
Soil Map Unit Name:				NIM/I Classification	
		2./:			on: Yes
Are climatic/hydrologic conditions on the site typical No No No No					
Are Vegetation, Soil, or Hydrology	naturally problem	natic? (If need	ed, explain any	answers in Remarks)	
SUMMARY OF FINDINGS - Attach site map showing		t locations, tra	nsects, importa	ant features, etc.	
Hydrophytic Vegetation Present?	lo —	Is the Sam	pled Area		
Hydric Soil Present?	lo 	within a W	etland?	No ———	_
N Wetland Hydrology Present?	lo	If yes, option	onal Wetland Si	te ID:	
Remarks: (Explain alternative procedures here or in	a separate report	.)		·	
The upland is a pasture dominated by a variety of gr	asses and forbs.				
VEGETATION Liso scientific names of plants					
VEGETATION - Use scientific names of plants.	Absolute			Bt	
Tree Stratum (Plot Size:)	% Cover	Dominant Species?	Indicator Status	Dominance Test worksheet: Number of Dominant Species	
rree stratum (Fiot Size)		speciesr	Status	That Are OBL, FACW, or FAC: 0	(4)
2				Total Number of Dominant	(A)
				2	
3				Species Across All Strata:	(B)
4				Percent of Dominant Species	
	0	= Total Cover		0 That Are OBL, FACW, or FAC:	(A/B)
Sapling/Shrub Stratum (Plot Size: 15)				Prevalence Index worksheet:	
1. Cornus alba	2.00	No	FACW	Total % Cover of:	Multiply by:
2				OBL species 0.00	x1 <u>0</u>
3				FACW species 2.00	x 2 <u>4</u>
4				FACU species 0.00	x 3 <u>420</u>
5				UPL species 15.00	x 4 <u>75</u>
	2	= Total Cover		Column Totals 122	(A) <u>499</u> (B)
Herb Stratum (Plot Size: 5)				Prevalence Index = B,	/A = 4.09016
1. Trifolium repens	45.00	Yes	FACU	Hydrophytic Vegetation Indicator	s:
2. Poa pratensis 3. Bromus inermis	35.00	Yes	FACU	1 - Rapid Test for Hydrop	, ,
Torque officinals	15.00	No	UPL	no 2 - Dominance Test is > 50	
Postulia alemente	15.00	No No	FACU	- S Trevalence mack is a	_
Thinonyrum intermedium	5.00	No No	FACU	4 - Morphological Adapta supporting data in Remarks or o	
0	3.00	140		Problematic Hydrophytic Vegetatio	_n 1
7 8.			-	(Explain)	""
0.	-			Indicators of hydric soil and wetland hydro	Now must be present
9				unless disturbed or problematic.	nogy must be present,
10					
	125	= Total Cover			
Woody Vine Stratum (Plot Size:)					
1.					
2.	-			1	
2				-	
	0	= Total Cover			
% Bare Ground in Herb Stratum 5				Hydrophytic Vegetation	
				Present?	
Remarks:					
Vegetation is dominated by Kentucky bluegrass and white clo	/er.				

SOIL Sampling Point: u-156n46...

inches) Color (moist) 0-9 10YR 2 1 9-13 2.5YR 5 6 13-18 10YR 3 1	%	C-1 (:-+)					
2.5YR 5 6		Color (moist)	%	Type ¹	Loc ²	Texture	Remarks
	100					sicl	
3 18 10VB 3 1	100					FS	fine sand
3-19 1011/2 1	100					LFS	loamy fine sand
							-
							-
ives: C-Concentration D-Depletion B	M-Roducod Matri	v MS=Macked Sand Gr					
ype: C=Concentration, D=Depletion, RI	vi=Reduced iviatri	x, MS=Masked Sand Gra	airis.			Indicator	rs for Problematic Hydric Soil ³ :
ydric Soil Indicators:		П					•
Histosol (A1)		Sandy Gleyed		4)			m Muck (A9) (LRR I, J)
Histic Epipedon (A2)		Sandy Redox					st Prairie Redox (A16)(LRR K, L, R)
Black Histic (A3)		Stripped Mati	rix (S6)			☐ Dar	k Surface (S7) (LRR G)
Hydrogen Sulfide (A4)		Loamy Mucky	Mineral (F1) (LRR I	K, L)	∐ Hig	h Plains Depressions (F16)
Stratified Layers (A5)		Loamy Gleyed	d Matrix (F	2)		(LRR	H outside of MLRA 72 & 73)
1cm Muck (A9) (LRR F, G, H)		Depleted Mat	trix (F3)			Rec	luced Vertic (F18)
Depleted Below Dark Surface (A11	.)	Redox Dark Si	urface (F6)		Rec	l Parent Material (F21)
Thick Dark Surface (A12)		Depleted Dark	k Surface	(F7)		☐ Ver	y Shallow Dark Surface (TF12)
¬		Redox Depres					ner (explain in remarks)
☐ Sandy Mucky Mineral (S1) ☐			, ,			011	ier (explain in remarks)
2.5cm Mucky Peat or Peat (S2)(LR		High Plains De					ors of hydrophytic vegetation and
5cm Mucky Peat or Peat (S3) (LRR	F)	(MLRA 72	& 73 of LI	RR H)			hydrology must be present, unless
						disturbe	d or problematic.
estrictive Layer (if present):							
Туре:		_				Hydric Soil Presen	t? No
Type: Depth (inches): marks: o hydric soil indicators were observed.						Hydric Soil Presen	t? <u>No</u>
Type: Depth (inches): emarks: Io hydric soil indicators were observed.						Hydric Soil Presen	t? <u>No</u>
	e is required; (:heck all that apply)					condary Indicators (minimum of two require
Type: Depth (inches): emarks: o hydric soil indicators were observed. YDROLOGY //etland Hydrology Indicators: rimary Indicators (minimum of or Surface Water (A1)	e is required; o	Salt Crust (B11)					condary Indicators (minimum of two require Surface Soil Cracks (B6)
Type: Depth (inches): emarks: o hydric soil indicators were observed. YDROLOGY //etland Hydrology Indicators: rimary Indicators (minimum of or Surface Water (A1) High Water Table (A2)	le is required; o	Salt Crust (B11) Aquatic Invertel	orates (B1				condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8)
Type: Depth (inches): emarks: o hydric soil indicators were observed. YDROLOGY //etland Hydrology Indicators: rimary Indicators (minimum of or Surface Water (A1) High Water Table (A2) Saturation (A3)	ue is required; o	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid	orates (B1 le Odor (C	1)			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10)
Type:	le is required; o	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wate	orates (B1 le Odor (C er Table (C	1)			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3)
Type:	ue is required; o	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wate	brates (B1 le Odor (C er Table (C pheres on	1)			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled)
Type:	ue is required; o	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wat Oxidized Rhizos (where not tilled	brates (B1 le Odor (C er Table (C pheres on	1) C2) Living Ro			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8)
Type:	e is required; o	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wat Oxidized Rhizos (where not tilled) Presence of Red	brates (B1 le Odor (C er Table (C pheres on) luced Iron	1) C2) Living Ro			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9)
Type:	e is required; o	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wate Oxidized Rhizos (where not tilled) Presence of Red Thin Muck Surfa	brates (B1 de Odor (C er Table (C pheres on) luced Iron ace (C7)	1) C2) Living Ro (C4)			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2)
Type:		Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wat Oxidized Rhizos (where not tilled) Presence of Red	brates (B1 de Odor (C er Table (C pheres on) luced Iron ace (C7)	1) C2) Living Ro (C4)			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Type:		Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wate Oxidized Rhizos (where not tilled) Presence of Red Thin Muck Surfa	brates (B1 de Odor (C er Table (C pheres on) luced Iron ace (C7)	1) C2) Living Ro (C4)			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2)
Type:		Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wate Oxidized Rhizos (where not tilled) Presence of Red Thin Muck Surfa	orates (B1 le Odor (C er Table (C pheres on) luced Iron ace (C7) n Remarks	1) C2) Living Ro (C4)			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Type:	ery (B7)	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wate Oxidized Rhizos (where not tilled) Presence of Red Thin Muck Surfa Other (Explain in	prates (B1 e Odor (C er Table (C pheres on) luced Iron ace (C7) n Remarks	1) C2) Living Ro (C4)			condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Type:	ery (B7)	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wate Oxidized Rhizos (where not tilled Presence of Red Thin Muck Surfa Other (Explain in	prates (B1 e Odor (C er Table (C pheres on) luced Iron ace (C7) n Remarks nes)	1) C2) Living Ro (C4)		Se	condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Type:	ery (B7) No No	Salt Crust (B11) Aquatic Invertel Hydrogen Sulfid Dry-Season Wate Oxidized Rhizos (where not tilled) Presence of Red Thin Muck Surfa Other (Explain in	prates (B1 e Odor (C er Table (C pheres on) luced Iron ace (C7) n Remarks nes)	1) C2) Living Ro (C4)		Se	condary Indicators (minimum of two require Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)