WETLAND DETERMINATION DATA FORM Great Plains Region

| Project/Site: Applicant: Investigators Soil Unit: | Enbridge | | | | | • | A or LRR): I Classification | MLRA 56 | | County: | 09/11/14 Pennington MN | | |
|---|---|---|--------|----------|---------------------------------------|---|--|---|--|----------------|---|---|--|
| Landform: | Depression | ession Local Relief: CL | | | | | | | | | Sample Point: | w-154n45w25-b1 | |
| Slope (%): | 0 - 2% Latitude: 48.13437483 Longitude: -96.3707226667 Datum: | | | | | | | | | - Constinue | | | |
| Are Vegetatio | • | I ⊠, or Hydrology | | | | al ? (If no, exp | 1 | ^{arks)} e normal circur | | □ No esent? | Section: Township: | | |
| Are Vegetatio | • | I □, or Hydrology | • | | | | | e normal circui ☑ Yes | | cocht: | Range: | Dir: | |
| SUMMARY C | | | | <i>J</i> | | | | | | | 5 | | |
| Hydrophytic \ | - | | | Yes | | - | | | | Is Present? | | | |
| Wetland Hyd | | | | Yes | depression in a highly disturbed area | | | | Is This Sampling Point Within A Wetland? Yes | | | | |
| Remarks: The wetland sample point is a wet meadow depression in a highly disturbed area. | | | | | | | | | | | | | |
| HYDROLOG | Y | | | | | | | | | | | | |
| | A1 - Surface A2 - High Wa A3 - Saturatic B1 - Water M B2 - Sedimer B3 - Drift Dep B4 - Algal Ma B5 - Iron Dep B7 - Inundatic | iter Table on arks at Deposits oosits t or Crust | | oly; Mi | | B11 - Salt B13 - Aqua C1 - Hydro C2 - Dry S C3 - Oxidiz | Crust atic Fauna ogen Sulfic eason Wa zed Rhizos ence of Re Muck Surfa | a de Odor ater Table spheres on Living educed Iron | | Secondary | B6 - Surface Soi B8 - Sparsely Ve B10 - Drainage I C3 - Oxidized RI C8 - Crayfish Bu C9 - Saturation V D2 - Geomorphic D5 - FAC-Neutra | egetated Concave Surface Patterns hizospheres on Living Roots (tilled) irrows Visible on Aerial Imagery c Position | |
| Field Observations: Surface Water Present? Yes Depth: (in.) Water Table Present? Yes Depth: 15 Saturation Present? Yes Depth: 7 Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available: Yes Remarks: The wetland sample point is located in a depression with a dry season water table and soil saturation observed. | | | | | | | | | | | | | |
| SOILS Profile Description (Describe to the depth needed to document the indicator or confirm the absence of indicators.) (Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered/Coated Sand Grains; Location: PL=Pore Lining, M=Matrix) | | | | | | | | | | | | | |
| | | , | , | | | | | J, | | | - | | |
| | Matrix | | | Mottl | | | | | | | | | |
| Depth (In.) | | Color (Moist) | | % | Color (I | Moist) | % | Туре | Location | Texture | | Remarks | |
| 0-3 | Hue_10YR | | | 100 | | 5/1 | 10 | | N A | SCL LFS | fine sand | | |
| <u>3-14</u> 14-21 | Hue_10YR Hue_10YR | | | 60 10 | Hue_10YR Hue_10YR | | 40 90 | D C | M | FS | | | |
| | | 171 | | 10 | | 0/0 | 50 | | | | | | |
| | | | | | | | | | | | | | |
| | | | | | | | | | | | | | |
| NRCS Hydr □ □ □ □ □ □ □ □ □ □ | A2 - Histic Epipedon A3 - Black Histic A4 - Hydrogen Sulfide A5 - Stratified Layers (LRR F) A9 - 1 cm Muck (LRR FGH) A11 - Depleted Below Dark Surface A12 - Thick Dark Surface S1 - Sandy Mucky Mineral S2 - 2.5 cm Mucky Peat or Peat (LRR F) S3 - 5 cm Mucky Peat or Peat (LRR F) S3 - 5 cm Mucky Peat or Peat (LRR F) S6 - Stripped Matrix S6 - Stripped Matrix S6 - Stripped Matrix F1 - Loamy Mucky Mineral F2 - Loamy Gleyed Matrix F3 - Depleted Matrix F6 - Redox Dark Surface F7 - Depleted Dark Surface F8 - Redox Depressions F16 - High Plains Depress | | | | | | | □ Indicators for Problematic Soils ¹ A9 - 1 cm Muck (LRR I, J) A16 - Coast Prairie Redox (LRR F, G, H) S7 - Dark Surface (LRR G) F16 - High Plains Depressions (LRR H, outside MLRA 72, 73) F18 - Reduced Vertic TF2 - Red Parent Material TF12 - Very Shallow Dark Surface Other (Explain in Remarks) (MLRA 72, 73 of LRR H) ¹ Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic. | | | | | |
| Restrictive Layer | Туре: | | Depth: | | | | Hydric Sc | Hydric Soil Present? Y | | | | | |
| Remarks: | - | Remarks: Soil is a dark layer of sandy clay loam underlain by a layer of loamy fine sand. The bottom layer is a light fine sand. Soil meets indicator A12 - Thick Dark Surface. | | | | | | | | | _ | | |

WETLAND DETERMINATION DATA FORM Great Plains Region

| Project/Site | : L3R | | | | Sample Point: w-154n45w25-b1 | | | | | |
|---|--|----------------|-----------------|------------|---|--|--|--|--|--|
| | | | | | | | | | | |
| VEGETATIC | | re non-native | species.) | | | | | | | |
| Tree Stratum | (Plot size: 30 ft. radius) | | | | Deminence Test Werkehest | | | | | |
| 4 | <u>Species Name</u> | <u>% Cover</u> | <u>Dominant</u> | Ind.Status | Dominance Test Worksheet | | | | | |
| 1. | | | | | Number of Deminent Oracias that are ODL EAOM as EAO (A) | | | | | |
| 2. | | | | | Number of Dominant Species that are OBL, FACW, or FAC: 4 (A) | | | | | |
| 3. | | | | | | | | | | |
| 4. | | | | | Total Number of Dominant Species Across All Strata: 4 (B) | | | | | |
| 5. | | | | | | | | | | |
| 6. | | | | | Percent of Dominant Species That Are OBL, FACW, or FAC: <u>100.0%</u> (A/B) | | | | | |
| 7. | | | | | | | | | | |
| 8. | | | | | Prevalence Index Worksheet | | | | | |
| 9. | | | | | Total % Cover of: Multiply by: | | | | | |
| 10. | | | | | OBL spp. 73 X 1 = 73 FACW spp. 30 X 2 = 60 FAC spp. 0 X 3 = 0 FACU spp. 0 X 4 = 0 | | | | | |
| | Total Cover = | 0 | | | FACW spp. 30 x 2 = 60 | | | | | |
| | | | | | FAC spp. 0 $x 3 = 0$ | | | | | |
| Sapling/Shrub | Stratum (Plot size: 15 ft. radius) | | | | FACU spp. 0 $x 4 = 0$ | | | | | |
| 1. | | | | | UPL spp. 0 $x 5 = 0$ | | | | | |
| 2. | | | | | | | | | | |
| 3. | | | | | Total <u>103</u> (A) <u>133</u> (B) | | | | | |
| 4. | | | | | | | | | | |
| 5. | | | | | Prevalence Index = B/A = 1.291 | | | | | |
| 6. | | | | | | | | | | |
| 7. | | | | | | | | | | |
| 8. | | | | | Hydrophytic Vegetation Indicators: | | | | | |
| 9. | | | | | Rapid Test for Hydrophytic Vegetation | | | | | |
| 10. | | | | | X Dominance Test is > 50% | | | | | |
| | | 0 | | | $\frac{1}{X} \qquad \text{Prevalence Index is } \leq 3.0 \text{ *}$ | | | | | |
| | | v | | | Morphological Adaptations (Explain) * | | | | | |
| Horb Stratum | (Plot pizer 5 ft rodiup) | | | | | | | | | |
| 1. | (Plot size: 5 ft. radius) Eleocharis obtusa | 25 | V | OBL | Problem Hydrophytic Vegetation (Explain) * | | | | | |
| | | | T V | | * Indicators of hydric soil and watland hydrology must be | | | | | |
| 2. | Beckmannia syzigachne | 20 | Y Y | OBL | * Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic. | | | | | |
| 3. | Juncus nodosus | 15 | · · · | OBL | · | | | | | |
| 4. | Juncus torreyi | 15 | Y | FACW | Definitions of Vegetation Strata: | | | | | |
| 5. | Juncus alpinoarticulatus | 10 | N | OBL | _ | | | | | |
| 6 | Argentina anserina | 5 | N | FACW | Tree - Woody plants 3 in. (7.6cm) or more in diameter at breast | | | | | |
| 7. | Hordeum jubatum | 5 | N | FACW | height (DBH), regardless of height. | | | | | |
| 8. | Asclepias incarnata | 3 | N | FACW | | | | | | |
| 9. | Lycopus americanus | 3 | N | OBL | Sapling/Shrub - Woody plants less than 3 in. DBH, regardless of height. | | | | | |
| 10. | Symphyotrichum lateriflorum | 2 | N | FACW | | | | | | |
| 11. | | | | | | | | | | |
| 12. | | | | | Herb - All herbaceous (non-woody) plants, regardless of size. | | | | | |
| 13. | | | | | | | | | | |
| 14. | | | | | | | | | | |
| 15. | | | | | Woody Vines - All woody vines, regardless of height. | | | | | |
| | Total Cover = | 103 | | | | | | | | |
| | | | | | | | | | | |
| Woody Vine S | tratum (Plot size: 30 ft. radius) | | | | | | | | | |
| 1. | | | | | | | | | | |
| 2. | | | | | | | | | | |
| 3. | - | | | | Hydrophytic Vegetation Present? Y | | | | | |
| 5. | <u>,</u> | | | | | | | | | |
| 4. | | | | | | | | | | |
| <u>т.</u> | Total Cover = | 0 | | | | | | | | |
| Remarks: The wetland sample point is dominated by blunt spike-rush, slough grass, knotted rush and Torrey's rush. | | | | | | | | | | |
| The marks. The wettand sample point is dominated by blunt spike-rush, slough grass, knotted rush and Torrey's rush. | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| Additional Remarks: | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |