## WETLAND DETERMINATION DATA FORM - Northcentral and Northeast Region

| Project/Site: L3R  | City/County: Clearwater                            | Sampling Date: 10/8/2014                              |
|--|--|---|
| Applicant/Owner: Enbridge  | State: MN  | 1 3   |
| Investigator(s): BJC/RAJ   |  | ownship, Range:                                       |
| Landform (hillslope, terrace, etc.): Footslope   |  | ncave, convex, none CL                                |
| Slope (%): 3 - 7% Lat.: 47.689517  | _Long.: <u>-95.394333</u> Datum                    |   |
| Soil Map Unit Name: 38B  Are climatic/hydrologic conditions of the site typical                            | for this time of the year?                         | NWI Classification:  (If no, explain in remarks)      |
| Are vegetation , soil , or hydrol  |  |   |
| Are vegetation , soil , or hydrol  |  | circumstances" present?                               |
| (If needed, explain any answers in remarks)  | natarany problematic.                              |   |
| ( · · · · · · · · · · · · · · · · · · ·  |  |   |
| SUMMARY OF FINDINGS  |  |   |
|  |  |   |
| Hydrophytic vegetation present?  | Is the sampled area withi                          | n a wetland? N  |
| Hydric soil present?  N  | -  | 10  |
| Indicators of wetland hydrology present?  N  | If yes, optional wetland site                      | e iD:   |
| Remarks: (Explain alternative procedures here or in a separate report.)                                    |  |   |
| The upland sample point is located in a hayfield dominated by red clover and Kentucky bluegrass.           |  |   |
| The apiana sample point is resulted in a may.  | iola dominatoa by roa olovor all                   | a Homaony Diaegrace.                                  |
|  |  |   |
|  |  |   |
| HYDROLOGY  |  |   |
|  |  | Secondary Indicators (minimum of two                  |
| Primary Indicators (minimum of one is required; che  |  | required)   |
|  | ater-Stained Leaves (B9)                           | Surface Soil Cracks (B6)                              |
|  | quatic Fauna (B13)                                 | Drainage Patterns (B10)                               |
|  | arl Deposits (B15)<br>ydrogen Sulfide Odor (C1)    | ☐ Moss Trim Lines (B16) ☐ Dry-Season Water Table (C2) |
|  | xidized Rhizospheres on Living                     | Crayfish Burrows (C8)                                 |
|  | oots (C3)  | Saturation Visible on Aerial Imagery                  |
|  | resence of Reduced Iron (C4)                       | (C9)  |
|  | ecent Iron Reduction in Tilled                     | Stunted or Stressed Plants (D1)                       |
|  | pils (C6)  | Geomorphic Position (D2)                              |
|  | nin Muck Surface (C7)<br>ther (Explain in Remarks) | Shallow Aquitard (D3)  Microtopographic Relief (D4)   |
| Sparsely Vegetated Concave   | iller (Explain in Remarks)                         | FAC-Neutral Test (D5)                                 |
| 34/1400 (50)   |  |   |
| Field Observations:  |  |   |
| Surface water present? Yes   | Depth (inches):                                    | _ Indicators of                                       |
| Water table present? Yes   | Depth (inches):                                    | wetland   |
| Saturation present? Yes  | Depth (inches):                                    | hydrology   |
| (includes capillary fringe)  |  | present? N  |
| Describe recorded data (stream gauge, monitoring well, aerial photos, previous inspections), if available: |  |   |
|  |  |   |
|  |  |   |
| Pamarks:   |  |   |
| Remarks: No indicators of wetland hydrology were of  |  |   |
| No indicators of wetland hydrology were observed.  |  |   |
| Two indicators of wetland flydrology were of   | oservea.   |   |