WETLAND DETERMINATION DATA FORM - North Central and Northeast Region

| Project/Site: SPP | City/County: Cass | | Sampling Date: 2016-08-03 | | | |
|---|--|------------------------------|--|--|--|--|
| Applicant/Owner: Enbridge | | State: Minnesota | Sampling Point: u-139n25w4-aa1 | | | |
| Investigator(s): DPT, MGH | Section, Towns | hip, Range: S4, T139N, R25 | | | | |
| Landform (hillslope, terrace, etc.): Side S | <u></u> | Local Relief (concave, co | | | | |
| Subregion (LRR or MLRA): | | • | gitude: -93.83826632 Datum: NAD83 | | | |
| Soil Map Unit Name: 146B | | | NWI Classification: N/A | | | |
| • | e site typical for this time of ve | ar? (if no explain in Remar | | | | |
| | | | | | | |
| Are Vegetation No_, Soil No_, or Hy | drology No significantly distu | urbed? Are "Normal Circum | nstances" present? Yes | | | |
| Are Vegetation No , Soil No , or Hydr | ology No naturally problema | atic? (If needed, explain an | y answers in Remarks) | | | |
| · — — , | · · | , , , | , | | | |
| SUMMARY OF FINDINGS - Attach site | map showing sampling point | locations, transects, impor | rtant features, etc. | | | |
| Hydrophytic Vegetation Present? | No | Is the Sampled Area | | | | |
| Hydric Soil Present? | No | within a Wetland? | <u>No</u> | | | |
| Wetland Hydrology Present? | <u>No</u> | If yes, optional Wetland | Site ID: | | | |
| Remarks: (Explain alternative procedure | es here or in a separate report. |) | | | | |
| No digging, existing road, potential buri | ied utilities. | | | | | |
| | | | | | | |
| | | | | | | |
| HYDROLOGY | | | | | | |
| i | | | Secondary Indicators (minimum of two required) | | | |
| Wetland Hydrology Indicators: Secondary Indicators (minimum of two required) | | | | | | |
| Primary Indicators (minimum of one is re | _ | (2.2) | Surface Soil Cracks (B6) | | | |
| Surface Water (A1) | Water-Stained Lea | | Drainage Patterns (B10) | | | |
| High Water Table (A2) | Aquatic Fauna (B1 | | Moss Trim Lines (B16) | | | |
| Saturation (A3) Water Marks (B1) | Marl Deposits (B1: Hydrogen Sulfide (| | Dry-Season Water Table (C2) Crayfish Burrows (C8) | | | |
| Sediment Deposits (B2) | | eres on Living Roots (C3) | Saturation Visible on Aerial Imagery (C9) | | | |
| Drift Deposits (B3) | Presence of Reduc | - | Stunted/Stressed Plants (D1) | | | |
| Algal Mat or Crust (B4) | | tion in Tilled Soils (C6) | Geomorphic Position (D2) | | | |
| Iron Deposits (B5) | Thin Muck Surface | | Shallow Aquitard (D3) | | | |
| Inundation Visible on Aerial Imagery (B7) | | | Microtopographic Relief (D4) | | | |
| Sparsely Vegetated Concave Surface (B8) | | | FAC-Neutral Test (D5) | | | |
| Field Observations: | | | | | | |
| Surface Water Present? | No Depth (inche | es) | | | | |
| Water Table Present? | Depth (inche | es) | | | | |
| Saturation Present? | No Depth (inche | es) | Wetland Hydrology Present? No | | | |
| (includes capillary fringe) | | | | | | |
| Describe Recorded Data (stream gauge, | monitoring well, aerial photos, | previous inspections), if av | railable: | | | |
| | | | | | | |
| Remarks: | | | | | | |
| No digging, could not verify water table | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| Ī. | | | | | | |

Tree Stratum

1. Populus tremuloides

(Plot Size: 30

| 2. Quercus palustris | 5.00 | No | FAC | Total Number of Dominant |
|---|---------|---------------|------|---|
| 3 | | | | Species Across All Strata: 5 (B) |
| 4. | | | | Percent of Dominant Species |
| 5. | | | | That Are OBL, FACW, or FAC: 20 (A/B) |
| 6. | | | | Prevalence Index worksheet: |
| 7. | | | | Total % Cover of: Multiply by: |
| | 55 | = Total Cover | | OBL species 0.00 x 1 0 |
| Sapling/Shrub Stratum (Plot Size: 15) | | | | FACW species 0.00 x 2 0 |
| 1. Corylus cornuta | 70.00 | Yes | UPL | FACU species 75.00 x 3 300 |
| 2. | | | | UPL species 70.00 x 4 350 |
| 3. | | - | | Column Totals 210 (A) 845 (B) |
| 4. | _ | | | Prevalence Index = B/A = 4.0238095 |
| 5. | | | | Hydrophytic Vegetation Indicators: |
| 6. | | | | 1 - Rapid Test for Hydrophytic Vegetation |
| 7. | | | | no 2 - Dominance Test is > 50% |
| ·· | 70 | = Total Cover | | no 3 - Prevalence Index is ≤ 3.0 ¹ |
| Herb Stratum (Plot Size: 5 | <u></u> | _ Total Cover | | 4 - Morphological Adaptations (Provide |
| 1. Eurybia macrophylla | 30.00 | Yes | FACU | supporting data in Remarks or on a separate sheet) |
| 2. Pteridium aquilinum | 25.00 | Yes | FACU | Problematic Hydrophytic Vegetation ¹ (Explain) |
| 3. Aralia nudicaulis | 20.00 | Yes | FACU | 1 |
| 4. Clintonia borealis | 10.00 | No | FAC | Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic. |
| 5. | | | | Definitions of Vegetation Strata: |
| 6. | | | | |
| 7. | | | | Tree - Woody plants 3 in. (.76 cm) or more in diameter at breast |
| 8. | | | | height (DBH), regardless of height. |
| 9 | | | | Sapling/Shrub - Woody plants less than 3 in. DBH and greater than |
| | | | | or equal to 3.28 ft (1 m) tall. |
| 10. | _ | | • | Herb - All herbaeceous (non-woody) plants, regardless of size, and |
| 11. | | | | woody plants less than 3.28 ft tall. |
| 12. | 85 | = Total Cover | - | Woody vines - All woody vines greater than 3.28 ft in height. |
| Woody Vine Stratum (Plot Size: 30) | | | | , |
| 1. | | | | |
| | | | | - Hydrophytic |
| 2 | | - | - | Vegetation |
| 3 | | - | | Present? |
| 4 | 0 | | | 1 |
| | | _=Total Cover | | |
| Remarks: (include photo numbers here or on a separate sheet | .) | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

Absolute

% Cover

50.00

Dominant

Species?

Yes

Indicator

Status

FAC

Sampling Point: u-139n25... **SOIL** Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.) Depth Matrix **Redox Features** Loc² (inches) Color (moist) Color (moist) % Type¹ Texture Remarks ¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, MS=Masked Sand Grains. ²Location: PL=Pore Lining, M=Matrix. Indicators for Problematic Hydric Soil³: Hydric Soil Indicators: Polyvalue Below Surface (S8) (LRR R, MLRA Histosol (A1) 2 cm Muck (A10) (LRR K, L, MLRA 149B) Histic Epipedon (A2) Coast Prairie Redox (A16)(LRR K, L, R) Thin Dark Surface (S9) (LRR R, MLRA 149B) Black Histic (A3) Loamy Mucky Mineral (F1) (LRR K, L) 5 cm Mucky Peat or Peat (S3) (LRR K, L, R) Hydrogen Sulfide (A4) Dark Surface (S7) (LRR K, M) Loamy Gleyed Matrix (F2) Stratified Layers (A5) Depleted Matrix (F3) Polyvalue Below Surface (S8) (LRR K, L) Depleted Below Dark Surface (A11) Redox Dark Surface (F6) Thin Dark Surface (S9) (LRR K, L) Thick Dark Surface (A12) Depleted Dark Surface (F7) Iron-Maganese Masses (F12) (LRR K, L, R) Sandy Mucky Mineral (S1) Redox Depressions (F8) Piedmont Floodplain Soils (F19) (MLRA 149B) Mesic Spodic (TA6) (MLRA 144A, 145, 149B) Sandy Gleyed Matrix (S4) Sandy Redox (S5) Red Parent Material (F21) Stripped Matrix (S6) Very Shallow Dark Surface (TF12) Dark Surface (S7) (LRR R, MLRA 149B) Other (explain in remarks) Restrictive Layer (if observed): Hydric Soil Present? No Depth (inches): Remarks: No digging, soils assumed non-hydric based on veg/hydro.

Site Photograph 1 Sampling Point: u-139n25w4-aa1



| Latitude: 46.8774456438114 | Cowardin Classification: | |
|------------------------------|--------------------------|--|
| Longitude: -93.8382652402798 | Circular 39: | |
| Direction: south | Eggers & Reed: | |
| Remarks: | | |
| Upland | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Site Photograph 2 Sampling Point: u-139n25w4-aa1



| Latitude: | 46.8774592224945 | Cowardin Classification: |
|-----------------|-------------------|--------------------------|
| Longitude: | -93.8382676710317 | Circular 39: |
| Direction: east | t | Eggers & Reed: |
| Remarks: | | |
| Upland | | |
| | | |
| | | |
| | | |
| | | |
| | | |